BENNETT

AI-POWERED BIOLOGICAL INTELLIGENCE

IDENTIFYN RESOLUTION STEPDOWN

SECONDARY ANTIBODY MICROSCOPY EVIDENCE ASSESSMENT

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse, F(ab')□ (H + L)

WESTERN BLOT -> WIDEFIELD -> WIDEFIELD SIM / APOTOME -> CONFOCAL -> AIRYSCAN
SIM -> SIM² -> SINGLE-MOLECULE STORM

8	1	PENDING
ORDERED EVIDENCE TIERS	SUPPORTING CONTROL	SCIENTIST REVIEW



REPRESENTATIVE HIGHEST-AVAILABLE VIEW / SINGLE-MOLECULE STORM

SA-000021 | PRODUCT-LEVEL EVIDENCE DOSSIER

SOURCE-GROUNDED MICROSCOPY + BIOLOGICAL REVIEW / IDENTIFYN EVIDENCE PRESERVED

CRO CAPABILITY DEMONSTRATED

Identifyn performed the experimental and imaging work documented below. BENNETT converts that operating history into a governed, source-bounded scientific record.

- Source-documented sample preparation and antibody/reagent workflow
- Western blot imaging
- Widefield fluorescence microscopy
- Widefield structured illumination / Apotome optical sectioning
- Confocal microscopy
- Airyscan acquisition and reconstruction workflow
- Structured illumination microscopy
- SIM² reconstruction workflow
- Single-molecule STORM presentation workflow
- Preserved control experiment
- Z-stack / 3D acquisition
- Multi-channel imaging

SCIENTIFIC QUESTION

What source-bounded experimental and microscopy evidence is preserved for Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse, F(ab')□ (H + L), and what is the strongest interpretation allowed by the available modalities and controls?

EVIDENCE AVAILABLE

Western blot -> Widefield -> Widefield SIM — Apotome -> Confocal -> Airyscan -> SIM -> SIM² -> Single-molecule STORM -> Control

BENNETT ASSESSMENT

The preserved SA-000021 record contains the full ordered evidence sequence from Western blot through single-molecule STORM. Each modality retains its own acquisition and processing boundary.

BENNETT / MICROSCOPY AND BIOLOGICAL EVIDENCE REVIEW

SOURCE-GROUNDED RETROSPECTIVE REVIEW / PENDING SCIENTIST REVIEW

SOURCE STATE	SOURCE HASH VERIFIED / EVENT RECORDED
BENNETT EVIDENCE REVIEW	COMPLETED / SOURCE-BOUNDED
NATIVE IMAGE REPROCESSING	NOT PERFORMED
LITERATURE SOURCES	INDEPENDENTLY VERIFIED
SCIENTIST REVIEW	PENDING
PUBLICATION	NOT AUTHORIZED

OVERALL CALL / FULL STEPDOWN / SOURCE-BOUNDED REVIEW

The preserved SA-000021 record contains the full ordered evidence sequence from Western blot through single-molecule STORM. Each modality retains its own acquisition and processing boundary.

SETTINGS ASSESSMENT

The record preserves 8 ordered evidence tier(s). Structured source metadata values are reported by stage and remain tied to the original method objects.

BIOLOGICAL SUPPORT

The record supports retrospective, source-bounded microscopy review. Target identity, specificity, treatment effect, binding, interaction and mechanism are not inferred from presentation images alone.

CONTROL ASSESSMENT

A true preserved control record is present and appears last.

CLAIM CEILING

This draft supports historical capability documentation, source registration, modality ordering, settings review and bounded interpretation. It does not independently validate antibody specificity, quantify a population response, establish binding, interaction, causation or mechanism, or authorize publication.

PROCESSING DISCLOSURE

Retrospective BENNETT architectural evaluation of preserved Identifyn source evidence. BENNETT did not perform native-file reprocessing, new deterministic measurement or the historical image post-processing represented in this record.

REAGENTS AND ACQUIRED CHANNELS

Preparation reagents are reported separately from channels actually documented in the acquisition settings. Fluorophores mentioned in staining are not automatically treated as acquired channels.

EVIDENCE OBJECT	REAGENTS / FLUOROPHORES PRESENT	CHANNELS ACTUALLY ACQUIRED
Western blot	488	Not deterministically stated in the acquisition block
Widefield	405/DAPI; 488; 594	2; 38 HE eGFP; 45 Texas Red; 450 - 490; 540 - 580; 500 - 550; 593 - 668; 493
Widefield SIM — Apotome	405/DAPI; 488; 594	2; 38 HE eGFP; 45 Texas Red; 450 - 490; 540 - 580; 500 - 550; 593 - 668; 493
Confocal	405/DAPI; 488; 594; 750	2; None; 488nm @0.2%; 561nm @0.2%; 493; 590; 517; 618
Airyscan	405/DAPI; 488; 594	2; None; 488nm @0.3%; 561nm @0.5%; 493; 590; 517; 618
SIM	405/DAPI; 488; 594	2; 488nm @3.50%; 561nm @4.0%; 488; 561; 655; 515; 488.000000
SIM²	405/DAPI; 488; 594	2; 488nm @3.50%; 561nm @4.0%; 488; 561; 655; 515; 488.000000
Single-molecule STORM	488	Both Channels
Control	405/DAPI; 488	2; 38 HE eGFP; 49 DAPI; 450 - 490; 335-383; 500 -550; 420-470; 493

MODALITY ASSESSMENT / PART 1 OF 9

Western blot / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: Azure Sapphire FL. Objective: not explicitly stated. Platform: Azure Sapphire FL with Customizable Laser and Filter Modules, Enabling UV- NIR and Z-Plane(s). Structured source metadata values: 11.

VISIBLE OBSERVATION

A preserved presentation image is available for Western blot. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Western blot, documents platform context as Azure Sapphire FL, and records detected dye/channel descriptors as 488.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy and antibody-validation guidance that presentation evidence must remain bounded by modality, controls and source documentation.

INTERPRETIVE LIMIT

Band position and displayed intensity do not independently establish analytical specificity, target identity, linearity, or treatment effect.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Western blot evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 2 OF 9

Widefield / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS Axio Imager.M1. Objective: Plan-Apochromat 63x/1.40 Oil M27. Platform: Zeiss Axio Imager.M1 Widefield, using a Plan-Apochromat 63x/1.40 Oil M27, Axiocam MR R3, and X-Cite Xylis Lamp was used with the following parameters for each dye:. Acquisition: Channels: 2; Scaling (Per Pixel): 0.102µm X 0.102µm; Image size (Pixels): 1202 X 925; Image Size (Pixels): 1202 X 925; Bit Depth: 12 Bit. Detector/camera: Imaging Device: Axiocam MR R3. Timing: Exposure Time: 200ms; 100ms. Illumination: Light source: X-Cite Xylis Lamp Intensity @40%; X-Cite Xylis Lamp Intensity @40.00%. Optical/scan: Effective NA: 1.4. Structured source metadata values: 30.

VISIBLE OBSERVATION

A preserved presentation image is available for Widefield. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Widefield, documents platform context as ZEISS Axio Imager.M1, and records detected dye/channel descriptors as 405/DAPI; 488; 594.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy and antibody-validation guidance that presentation evidence must remain bounded by modality, controls and source documentation.

INTERPRETIVE LIMIT

A presentation image supports gross localization and source-bounded co-occurrence, not direct binding, molecular colocalization, or native quantitative measurement.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Widefield evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 3 OF 9

Widefield SIM — Apotome / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS Axio Observer.Z1/7 Apotome. Objective: EC Plan-Neofluar 63X/1.25 oil M27. Platform: Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:. Acquisition: Channels: 2; Scaling (Per Pixel): 0.143µm X 0.143µm; Image size (Pixels): 1106 X 834; Image Size (Pixels): 1106 X 834; Bit Depth: 14 Bit. Detector/camera: Imaging Device: Axiocam 807. Timing: Exposure Time: 200ms; 100ms. Illumination: Light source: X-Cite Xylis Lamp Intensity @50%; X-Cite Xylis Lamp Intensity @25%. Optical/scan: Effective NA: 1.25. Structured source metadata values: 32.

VISIBLE OBSERVATION

A preserved presentation image is available for Widefield SIM — Apotome. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Widefield SIM — Apotome, documents platform context as ZEISS Axio Observer.Z1/7 Apotome, and records detected dye/channel descriptors as 405/DAPI; 488; 594.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy and antibody-validation guidance that presentation evidence must remain bounded by modality, controls and source documentation.

INTERPRETIVE LIMIT

Improved optical sectioning does not by itself establish reconstructed super-resolution or molecular interaction.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Widefield SIM — Apotome evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 4 OF 9

Confocal / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS LSM 900. Objective: with a Plan-Apochromat 63X/1.40 oil DIC M27. Platform: Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:. Acquisition: Channels: 2; Scaling (Per Pixel): 85.08nm X 85.08nm; Image size (Pixels): 1192 X 1192; Image Size (Pixels): 1192 X 1192; Bit Depth: 16 Bit. Detector/camera: Imaging Device: Airyscan; GaAsp-PMT 2; Detector Type: GaAsp-PMT. Timing: Pixel Time: 0.65µs; Frame Time: 8.67s. Illumination: Laser Wavelength: 488nm @0.2%; 561nm @0.2%. Optical/scan: Pinhole: 1.00AU / 45µm; 1.00AU / 54µm; Scan Zoom: 1.0; LSM Scan Speed: 8; Scan Direction: Bidirectional; Effective NA: 1.4. Structured source metadata values: 39.

VISIBLE OBSERVATION

A preserved presentation image is available for Confocal. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Confocal, documents platform context as ZEISS LSM 900, and records detected dye/channel descriptors as 405/DAPI; 488; 594; 750.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy and antibody-validation guidance that presentation evidence must remain bounded by modality, controls and source documentation.

INTERPRETIVE LIMIT

Optical sectioning and source-bounded 3D presentation do not establish binding, mechanism, or population-level effect.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Confocal evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 5 OF 9

Airyscan / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS LSM 900. Objective: with a Plan-Apochromat 63X/1.40 oil DIC M27. Platform: Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:. Acquisition: Z-Stack: 160 Slices (6.36µm); Channels: 2; Scaling (Per Pixel): 42.52nm X 42.52nmµm X 40.00nm; Image size (Pixels): 1535 X 1348; Image Size (Pixels): 1535 X 1348; Bit Depth: 16 Bit. Detector/camera: Imaging Device: Airyscan; Detector Type: GaAsp-PMT. Timing: Pixel Time: 0.66µs; Frame Time: 3.94s. Illumination: Laser Wavelength: 488nm @0.3%; 561nm @0.5%. Optical/scan: Pinhole: 5.31AU / 236µm; 5.00AU / 272µm; Scan Zoom: 1.5; LSM Scan Speed: 7; Scan Direction: Bidirectional; Effective NA: 1.4. Processing: Projection: View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected); Background: Black; AiryScan Mode: Super Resolution 9.1 (3D, Auto); Super Resolution 8.3 (3D, Auto). Structured source metadata values: 46.

VISIBLE OBSERVATION

A preserved presentation image is available for Airyscan. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Airyscan, documents platform context as ZEISS LSM 900, and records detected dye/channel descriptors as 405/DAPI; 488; 594.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy guidance that advanced-resolution outputs remain dependent on acquisition, reconstruction or localization quality control.

INTERPRETIVE LIMIT

Displayed feature separation remains reconstruction dependent; no unsupported numerical resolution or molecular architecture claim is permitted.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Airyscan evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 6 OF 9

SIM / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS Elyra 7. Objective: Plan-Apochromat 63X/1.46 oil Korr M27. Platform: Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:. Acquisition: Z-Stack: 35Slices (2.04µm); Channels: 2; Scaling (Per Pixel): 31.30nm X 31.30nm X 60.00nm; Image size (Pixels): 2560 X 2560; Image Size (Pixels): 2560 X 2560; Bit Depth: 16 Bit. Detector/camera: Imaging Device: Edge 4.2 SIM; Detector Type: Camera. Timing: Exposure Time: 65ms; 60ms; Frame Time: 1.00s. Illumination: Laser Wavelength: 488nm @3.50%; 561nm @4.0%. Optical/scan: Scan Zoom: 1.0; Scan Direction: Unidirectional; Averaging: 8; Effective NA: 1.46. Processing: Projection: View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected); Background: Black. Structured source metadata values: 57.

VISIBLE OBSERVATION

A preserved presentation image is available for SIM. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as SIM, documents platform context as ZEISS Elyra 7, and records detected dye/channel descriptors as 405/DAPI; 488; 594.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy guidance that advanced-resolution outputs remain dependent on acquisition, reconstruction or localization quality control.

INTERPRETIVE LIMIT

Reconstruction-dependent structural separation requires native reconstruction QC before numerical resolution or molecular dimensions can be claimed.

CLAIM CEILING

This tier supports a source-bounded statement that the historical SIM evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 7 OF 9

SIM² / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS Elyra 7. Objective: Plan-Apochromat 63X/1.46 oil Korr M27. Platform: Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:. Acquisition: Z-Stack: 26Slices (1.5µm); Channels: 2; Scaling (Per Pixel): 15.65nm X 15.65nm X 60.00nm; Image size (Pixels): 4734 X 2971; 312 X 312; Image Size (Pixels): 4734 X 2971; 312 X 312; Bit Depth: 16 Bit. Detector/camera: Imaging Device: Edge 4.2 SIM; Detector Type: Camera. Timing: Exposure Time: 65ms; 60ms; Frame Time: 1.0s. Illumination: Laser Wavelength: 488nm @3.50%; 561nm @4.0%. Optical/scan: Scan Zoom: 1.0; Scan Direction: Unidirectional; Averaging: 8; Effective NA: 1.46. Processing: Projection: View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected); Background: Black. Structured source metadata values: 62.

VISIBLE OBSERVATION

A preserved presentation image is available for SIM². BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as SIM², documents platform context as ZEISS Elyra 7, and records detected dye/channel descriptors as 405/DAPI; 488; 594.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy guidance that advanced-resolution outputs remain dependent on acquisition, reconstruction or localization quality control.

INTERPRETIVE LIMIT

Further reconstruction-dependent segmentation does not establish molecular dimensions, structure identity, or mechanism without independent support.

CLAIM CEILING

This tier supports a source-bounded statement that the historical SIM² evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 8 OF 9

Single-molecule STORM / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: Bruker Vutara VXL. Objective: Olympus 60X/1.30 silicon oil objective. Platform: Bruker Vutara VXL Biplane Single Molecule Microscope, using an Olympus 60X/1.30 silicon oil objective, was used with the following parameters:. Detector/camera: Camera: Both Cameras. Timing: Exposure Time: 20ms Selected; Super Resolution Camera Exposure Time: 20ms; Widefield Camera Exposure Time: 100ms. Illumination: Photoactivation/Imaging Laser: 0% Selected. Structured source metadata values: 71.

VISIBLE OBSERVATION

A preserved presentation image is available for Single-molecule STORM. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Single-molecule STORM, documents platform context as Bruker Vutara VXL, and records detected dye/channel descriptors as 488.

LITERATURE CORRELATION

Consistent with peer-reviewed microscopy guidance that advanced-resolution outputs remain dependent on acquisition, reconstruction or localization quality control.

INTERPRETIVE LIMIT

Without localization tables, uncertainty, blinking, drift, and independent cluster evidence, the presentation cannot establish molecule count, cluster truth, exact precision, or molecular architecture.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Single-molecule STORM evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

MODALITY ASSESSMENT / PART 9 OF 9

Control / SOURCE EVIDENCE PRESERVED

DOCUMENTED SETTINGS

Platform: ZEISS Axio Imager.M1; ZEISS Axio Observer.Z1/7 Apotome. Objective: EC Plan-Neofluar 63X/1.25 oil M27. Platform: Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:. Acquisition: Channels: 2; Scaling (Per Pixel): 0.143µm X 0.143µm; Image size (Pixels): 1600 X 1104; Image Size (Pixels): 1600 X 1104; Bit Depth: 14 Bit. Detector/camera: Imaging Device: Axiocam 807 mono. Timing: Exposure Time: 200ms; 30ms. Illumination: Light source: X-Cite Xylis Lamp Intensity @25%; X-Cite Xylis Lamp Intensity @10%. Optical/scan: Effective NA: 1.25. Structured source metadata values: 30.

VISIBLE OBSERVATION

A preserved presentation image is available for Control. BENNETT records the visible source presentation without creating a new native measurement.

SOURCE-REPORTED RESULT

The source identifies this object as Control, documents platform context as ZEISS Axio Imager.M1; ZEISS Axio Observer.Z1/7 Apotome, and records detected dye/channel descriptors as 405/DAPI; 488.

LITERATURE CORRELATION

Consistent with peer-reviewed antibody-control guidance that a no-primary or related control bounds background but is not a complete specificity system.

INTERPRETIVE LIMIT

A control bounds the recorded comparison only; it does not establish universal specificity, zero background, zero cross-reactivity, or absence of free dye.

CLAIM CEILING

This tier supports a source-bounded statement that the historical Control evidence and documented settings are preserved for review; stronger biological or quantitative conclusions require compatible native data, controls and scientist review.

CROSS-MODALITY SYNTHESIS

The preserved SA-000021 record contains the full ordered evidence sequence from Western blot through single-molecule STORM. Each modality retains its own acquisition and processing boundary.

Different acquisitions are not treated as a quantitative intensity ladder. Any continuity statement remains qualitative and source bounded.

BENNETT LITERATURE CORRELATION — SOURCES INDEPENDENTLY VERIFIED

BENNETT uses independently verified primary scientific sources to test whether the interpretation is scientifically consistent and properly bounded. Literature correlation does not retroactively validate the Identifyn dataset or replace scientist review.

- Antibody and microscopy evidence require orthogonal controls and modality-appropriate interpretation; presentation images alone do not establish analytical specificity.

- Advanced-resolution presentation depends on acquisition, reconstruction or localization quality control before numerical resolution, molecular dimensions or cluster identity can be claimed.

REFERENCES

1. Uhlen M, et al. A proposal for validation of antibodies. Nat Methods. 2016;13:823-827. doi:10.1038/nmeth.3995.
2. Gustafsson MGL. Surpassing the lateral resolution limit by a factor of two using structured illumination microscopy. J Microsc. 2000;198:82-87. doi:10.1046/j.1365-2818.2000.00710.x.
3. Sage D, et al. Quantitative evaluation of software packages for single-molecule localization microscopy. Nat Methods. 2015;12:717-724. doi:10.1038/nmeth.3442.

RELEASE BOUNDARY

This assessment remains a draft pending scientist review. No publication is authorized. Any future native-file analysis must enter BENNETT as a new evidence snapshot with routed engine authority, native-data QA, methods, limitations, and an independently reviewed claim ceiling.

SOURCE RECONCILIATION / QA / PROVENANCE

SOURCE STATUS - PRESERVED / AUTO-RECONCILED

Historical Identifyn source material was subjected to BENNETT retrospective QA. Original source state and correction history remain preserved in provenance.

HIGH CONFIDENCE - AUTO-RECONCILED

Confocal / Scaling (Per Pixel): 85.08nm X 85.08nm -> 0.08508 µm X 0.08508 µm. Both numerical magnitude and unit were normalized, then checked against the reported field dimensions; no unit label was changed alone. Supporting evidence: Image Size (Pixels)=1192 X 1192; Image Size (Scaled)=101.41µm X 101.41µm; PASS - INTERNALLY CONSISTENT. Documented sampling reproduces the reported physical field dimensions. Maximum field-dimension error 0.01%.

HIGH CONFIDENCE - AUTO-RECONCILED

Airyscan / Scaling (Per Pixel): 42.52nm X 42.52nmµm X 40.00nm -> 0.04252 µm X 0.04252 µm X 0.04000 µm. Both numerical magnitude and unit were normalized, then checked against the reported field dimensions; no unit label was changed alone. Supporting evidence: Image Size (Pixels)=1535 X 1348; Image Size (Scaled)=65.27µm X 57.32µm; PASS - INTERNALLY CONSISTENT. Documented sampling reproduces the reported physical field dimensions. Maximum field-dimension error 0.01%.

HIGH CONFIDENCE - AUTO-RECONCILED

SIM / Scaling (Per Pixel): 31.30nm X 31.30nm X 60.00nm -> 0.03130 µm X 0.03130 µm X 0.06000 µm. Both numerical magnitude and unit were normalized, then checked against the reported field dimensions; no unit label was changed alone. Supporting evidence: Image Size (Pixels)=2560 X 2560; Image Size (Scaled)=80.14µm X 80.14µm; PASS - INTERNALLY CONSISTENT. Documented sampling reproduces the reported physical field dimensions. Maximum field-dimension error 0.01%.

HIGH CONFIDENCE - AUTO-RECONCILED

SIM² / Scaling (Per Pixel): 15.65nm X 15.65nm X 60.00nm -> 0.01565 µm X 0.01565 µm X 0.06000 µm. Both numerical magnitude and unit were normalized, then checked against the reported field dimensions; no unit label was changed alone. Supporting evidence: Image Size (Pixels)=4734 X 2971; Image Size (Scaled)=74.09µm X 46.50µm; PASS - INTERNALLY CONSISTENT. Documented sampling reproduces the reported physical field dimensions. Maximum field-dimension error 0.01%.

BENNETT

AI-POWERED BIOLOGICAL INTELLIGENCE

BENNETT EVIDENCE REVIEW COMPLETE

INTERPRETATION TO EVIDENCE

BENNETT has completed the bounded retrospective assessment.
The following pages switch ownership to the preserved Identifyn record.

IDENTiFYN®

IDENTIFYN PRESERVED SOURCE EVIDENCE

ORIGINAL RECORD BEGINS

Images, source methods, acquisition settings and provenance follow in the locked Stepdown order. Scientific wording is preserved; missing modalities are not invented.
A preserved control follows all available Stepdown tiers as supporting evidence.

PRODUCT-LEVEL STEPDOWN MAP

This dossier preserves the complete available evidence progression for SA-000021. Each tier remains tied to its original image, source method, instrument context and cryptographic identity. A preserved control is retained as supporting evidence and appears after every resolution tier.

CANONICAL STEPDOWN TIER	EVIDENCE STAGE	INSTRUMENT	STATUS
1	Western blot	Azure Sapphire FL	PRESERVED
2	Widefield	ZEISS Axio Imager.M1	PRESERVED
3	Widefield SIM - Apotome	ZEISS Axio Observer.Z1/7 Apotome	PRESERVED
4	Confocal	ZEISS LSM 900	PRESERVED
5	Airyscan	ZEISS LSM 900	PRESERVED
6	SIM	ZEISS Elyra 7	PRESERVED
7	SIM²	ZEISS Elyra 7	PRESERVED
8	Single-molecule STORM	Bruker Vutara VXL	PRESERVED

INTERPRETIVE BOUNDARY

Ordering evidence by resolution tier does not make the modalities interchangeable. Each stage retains its acquisition environment and scientific limitations. This pilot organizes the record; it does not synthesize new biological claims.

ORIGINAL IMAGE EVIDENCE / WESTERN BLOT



EVIDENCE TIER	Western blot
INSTRUMENT	Azure Sapphire FL
CELL MODEL	HeLa
DETECTED DYES	488
METADATA VALUES	11

STRUCTURED ACQUISITION METADATA / WESTERN BLOT

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Azure Sapphire FL with Customizable Laser and Filter Modules, Enabling UV- NIR and Z-Plane(s)
Scanner	Azure Sapphire FL with Customizable Laser and Filter Modules, Enabling UV- NIR and Z-Plane(s)
Detection mode	Fluorescence
Laser	488nm
Emission Filter	513±17nm
Detector	PMT
Intensity	2
Pixel size	100 µm
Scan Speed	High
Quality	High
Focus	Membrane (0 µm in Z-axis)
Image	Identifyn® LLC

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

HeLa cells were homogenized using an ultrasonication probe and centrifuged to pellet down the cell debris. The lysate supernatant was mixed with NuPAGE™ LDS Sample Buffer (4X), heated at 80oC for 10 min, and subjected to electrophoresis using NuPAGE™ 4 to 12%, Bis-Tris, 1.0-1.5 mm, Mini Protein Gels in a Mini Gel Tank. The Thermo Fisher's PageRuler™ Prestained Protein Ladder, 10 to 180 kDa, was used as a molecular weight marker.

Post electrophoresis, the proteins were transferred to a polyvinylidene difluoride (PVDF) membrane using iBlot™ 3 Starter Kit, PVDF. The transfer membrane was blocked with Blocker™ Casein for 1 hour and washed 5X with PBS. The blocked membrane was incubated with Actin Monoclonal Antibody (mAbGEa) (MA1-744) for 2 hours, followed by 5X PBS washes. Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit anti-Mouse F(ab)2 was incubated for 1 hour and washed 5X with PBS. The membrane was dried for 10 minutes at room temperature before imaging.

Scanner = Azure Sapphire FL with Customizable Laser and Filter Modules, Enabling UV- NIR and Z-Plane(s)

Scan Settings

Detection mode = Fluorescence
Laser = 488nm
Emission Filter = 513±17nm
Detector = PMT
Intensity = 2
Pixel size = 100 µm
Scan Speed = High
Quality = High
Focus = Membrane (0 µm in Z-axis)

Post Processing

None

Image: Identifyn® LLC

Copyright 2026 GMD12 LLC.

SOURCE PROVENANCE

Catalog	SA-000021
Stage	Western blot

SOURCE PROVENANCE

Instrument	Azure Sapphire FL
Microscope configuration	Azure Sapphire FL with Customizable Laser and Filter Modules, Enabling UV- NIR and Z-Plane(s)
Structured source metadata values	11
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	30EB5C3D3E3D495FD5BBC77D6D0E67C155206EF6C5F839E38F53F6DA955BFBAA
Method PDF SHA-256	D083623DD95BE3A5939ADF4AA1951E903B144BE5757EC4CA703DB1A643767B24

ORIGINAL IMAGE EVIDENCE / WIDEFIELD



EVIDENCE TIER	Widefield
INSTRUMENT	ZEISS Axio Imager.M1
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488; 594
METADATA VALUES	30

STRUCTURED ACQUISITION METADATA / WIDEFIELD

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss Axio Imager.M1 Widefield, using a Plan-Apochromat 63x/1.40 Oil M27, Axiocam MR R3, and X-Cite Xylis Lamp was used with the following parameters for each dye:
Objective	Plan-Apochromat 63x/1.40 Oil M27
Channels	2
Scaling (Per Pixel)	0.102µm X 0.102µm
Image size (Pixels)	1202 X 925
Image Size (Scaled)	123.06µm X 94.07µm
Bit Depth	12 Bit
Image Center Position	X: 0.00µm, Y: 0.00µm
ROI Center Offset	X: 0.00µm, Y: 0.00µm
Reflector	38 HE eGFP | 45 Texas Red
Beam Splitter	495 | 585
Filter Excitation Wavelength	450 - 490 | 540 - 580
Filter Emission Wavelength	500 - 550 | 593 - 668
Contrast Method	Fluorescence
Light source	X-Cite Xylis Lamp Intensity @40% | X-Cite Xylis Lamp Intensity @40.00%
Exposure Time	200ms | 100ms
Excitation Wavelength	493 | 590
Emission Wavelength	517 | 618
Effective NA	1.4
Imaging Device	Axiocam MR R3
Camera Adapter	1X
Depth of Focus	0.80µm | 0.96µm
Binning Mode	1,1

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

Identifyn® SRM Alexa Fluor™ 594 Affinity Purified Donkey Anti-Rabbit F(ab')□ (H + L)
Cat ID# - SA 000054

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher, Fish Serum Blocker cat#37527X3. The first primary antibody, Mitochondria Antibody (MA5-12017), was incubated overnight at 4°C at a dilution of 1:100. Cells were washed 3X in PBS. Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000.

The second primary antibody followed this, Actin Recombinant Rabbit Monoclonal Antibody (JF47-01), which was incubated for 1 hour at room temperature at a dilution of 1:250. Cells were washed 3X in PBS. Identifyn® 594 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss Axio Imager.M1 Widefield, using a Plan-Apochromat 63x/1.40 Oil M27, Axiocam MR R3, and X-Cite Xylis Lamp was used with the following parameters for each dye:

Single Plane (2D)

Channels = 2
Scaling (Per Pixel) = 0.102µm X 0.102µm
Image size (Pixels) = 1202 X 925
Image Size (Scaled) = 123.06µm X 94.07µm
Bit Depth = 12 Bit
Image Center Position = X: 0.00µm, Y: 0.00µm
ROI Center Offset = X: 0.00µm, Y: 0.00µm

488 -

Reflector = 38 HE eGFP
Beam Splitter = 495
Filter Excitation Wavelength = 450 - 490
Filter Emission Wavelength = 500 - 550
Contrast Method = Fluorescence
Light source = X-Cite Xylis Lamp Intensity @40%
Exposure Time = 200ms
Excitation Wavelength = 493
Emission Wavelength = 517
Effective NA = 1.4
Imaging Device = Axiocam MR R3
Camera Adapter = 1X
Depth of Focus = 0.80µm
Binning Mode = 1,1

594 -

Reflector = 45 Texas Red
Beam Splitter = 585
Filter Excitation Wavelength = 540 - 580
Filter Emission Wavelength = 593 - 668
Contrast Method = Fluorescence
Light source = X-Cite Xylis Lamp Intensity @40.00%
Exposure Time = 100ms
Excitation Wavelength = 590
Emission Wavelength = 618
Effective NA = 1.4
Imaging Device = Axiocam MR R3
Camera Adapter = 1X
Depth of Focus = 0.96µm
Binning Mode = 1,1

Post Processing

None

Image Correction

None

Image by E.F. Simon

SOURCE PROVENANCE

Catalog	SA-000021
Stage	Widefield
Instrument	ZEISS Axio Imager.M1
Microscope configuration	Zeiss Axio Imager.M1 Widefield, using a Plan-Apochromat 63x/1.40 Oil M27, Axiocam MR R3, and X-Cite Xylis Lamp was used with the following parameters for each dye:
Structured source metadata values	30
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	63C45F3ADD9FC97F286593218C01D5A0138D8E285082D092587BFCB52004CD34
Method PDF SHA-256	3EBC3247447AB27B4C53A595396AC47F04707C1F7392D6672DFD0E90AF4DBFD2

ORIGINAL IMAGE EVIDENCE / WIDEFIELD SIM - APOTOME



EVIDENCE TIER	Widefield SIM - Apotome
INSTRUMENT	ZEISS Axio Observer.Z1/7 Apotome
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488; 594
METADATA VALUES	32

STRUCTURED ACQUISITION METADATA / WIDEFIELD SIM - APOTOME

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:
Objective	EC Plan-Neofluar 63X/1.25 oil M27
Channels	2
Scaling (Per Pixel)	0.143µm X 0.143µm
Image Size (Pixels)	1106 X 834
Image Size (Scaled)	158.00µm X 119.14µm
Bit Depth	14 Bit
Image Center Position	X: -12.20mm, Y: -1.09mm
Stage Position	X: -12.20mm, Y: -1.09mm
ROI Center Offset	X: 1.14µm, Y: 0.00µm
Reflector	38 HE eGFP | 45 Texas Red
Beam Splitter	495 | 585
Filter Excitation Wavelength	450 - 490 | 540 - 580
Filter Emission Wavelength	500 - 550 | 593 - 668
Contrast Method	Fluorescence
Light source	X-Cite Xylis Lamp Intensity @50% | X-Cite Xylis Lamp Intensity @25%
Exposure Time	200ms | 100ms
Excitation Wavelength	493 | 590
Emission Wavelength	517 | 618
Effective NA	1.25
Imaging Device	Axiocam 807
Camera Adapter	1X
Depth of Focus	1.00µm | 1.07µm
Binning Mode	2,2
Apotome SIM Image Processing	Display Mode “Optical Sectioning” - enabled correction selected.

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

Identifyn® SRM Alexa Fluor™ 594 Affinity Purified Donkey Anti-Rabbit F(ab')□ (H + L)
Cat ID# - SA 000054

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher, Fish Serum Blocker cat#37527X3. The first primary antibody, Mitochondria Antibody (MA5-12017), was incubated overnight at 4°C at a dilution of 1:100. Cells were washed 3X in PBS. Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000.

The second primary antibody followed this, Actin Recombinant Rabbit Monoclonal Antibody (JF47-01), which was incubated for 1 hour at room temperature at a dilution of 1:250. Cells were washed 3X in PBS. Identifyn® 594 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:

Single Plane (2D)

Channels = 2
Scaling (Per Pixel) = 0.143µm X 0.143µm
Image Size (Pixels) = 1106 X 834
Image Size (Scaled) = 158.00µm X 119.14µm
Bit Depth = 14 Bit
Image Center Position = X: -12.20mm, Y: -1.09mm
Stage Position = X: -12.20mm, Y: -1.09mm
ROI Center Offset = X: 1.14µm, Y: 0.00µm

488 -

Reflector = 38 HE eGFP
Beam Splitter = 495
Filter Excitation Wavelength = 450 - 490
Filter Emission Wavelength = 500 - 550
Contrast Method = Fluorescence
Light source = X-Cite Xylis Lamp Intensity @50%
Exposure Time = 200ms
Excitation Wavelength = 493
Emission Wavelength = 517
Effective NA = 1.25
Imaging Device = Axiocam 807
Camera Adapter = 1X
Depth of Focus = 1.00µm
Binning Mode = 2,2

594 -

Reflector = 45 Texas Red
Beam Splitter = 585
Filter Excitation Wavelength = 540 - 580
Filter Emission Wavelength = 593 - 668
Contrast Method = Fluorescence
Light source = X-Cite Xylis Lamp Intensity @25%
Exposure Time = 100ms
Excitation Wavelength = 590
Emission Wavelength = 618
Effective NA = 1.25
Imaging Device = Axiocam 807
Camera Adapter = 1X
Depth of Focus = 1.07µm
Binning Mode = 2,2

Post Processing - SIM Processing

Apotome SIM Image Processing = Display Mode “Optical Sectioning” - enabled correction selected.

Normalization, Phase Correction, Fourier Filter, and Deconvolution were NOT used.

Image Corrections

None

Image by E.F. Simon

SOURCE PROVENANCE

Catalog	SA-000021
Stage	Widefield SIM - Apotome
Instrument	ZEISS Axio Observer.Z1/7 Apotome
Microscope configuration	Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:
Structured source metadata values	32
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	8B1EAB57011E3CBE222AA6376C517FCD00BFEE04051E759F88357AFC7028B054
Method PDF SHA-256	29D255819C4CD61CC9499088917234B8F9EFAAC4D438A539F21BEFD1E32BABB1

ORIGINAL IMAGE EVIDENCE / CONFOCAL



EVIDENCE TIER	Confocal
INSTRUMENT	ZEISS LSM 900
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488; 594; 750
METADATA VALUES	39

STRUCTURED ACQUISITION METADATA / CONFOCAL

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:
Objective	with a Plan-Apochromat 63X/1.40 oil DIC M27
Channels	2
Scaling (Per Pixel)	0.08508 µm X 0.08508 µm
Image Size (Pixels)	1192 X 1192
Image Size (Scaled)	101.41µm X 101.41µm
Bit Depth	16 Bit
Image Center Position	X: 72.64mm, Y: 50.012mm
Stage Position	X: 72.64mm, Y: 50.012mm
ROI Center Offset	X: 0.00µm, Y: 0.00µm
Reflector	None
Contrast Method	Fluorescence
Pinhole	1.00AU / 45µm | 1.00AU / 54µm
LMS MBS	MBS 405 + 488 + 561 + 640(T10/R90)
Laser Wavelength	488nm @0.2% | 561nm @0.2%
Laser Blanking	Enabled
Scan Mode	Frame
Scan Zoom	1.0
Rotation	0°
Sampling	1.00
Pixel Time	0.65µs
Frame Time	8.67s
LSM Scan Speed	8
Scan Direction	Bidirectional
Line Step	1
Excitation Wavelength	493 | 590
Emission Wavelength	517 | 618
Effective NA	1.4
Detection Wavelength	496 - 575 | 590 - 700
Imaging Device	Airyscan | GaAsp-PMT 2
Detector Type	GaAsp-PMT
Detector Gain	750 V | 650 V
Detector Offset	0

FIELD	SOURCE-DOCUMENTED VALUE
Detector Digital Gain	1.0

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

Identifyn® SRM Alexa Fluor™ 594 Affinity Purified Donkey Anti-Rabbit F(ab')□ (H + L)
Cat ID# - SA 000054

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher, Fish Serum Blocker cat#37527X3. The first primary antibody, Mitochondria Antibody (MA5-12017), was incubated overnight at 4°C at a dilution of 1:100. Cells were washed 3X in PBS. Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000.

The second primary antibody followed this, Actin Recombinant Rabbit Monoclonal Antibody (JF47-01), which was incubated for 1 hour at room temperature at a dilution of 1:250. Cells were washed 3X in PBS. Identifyn® 594 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:

Single Plane (2D)

Channels = 2
Scaling (Per Pixel) = 85.08nm X 85.08nm
Image Size (Pixels) = 1192 X 1192
Image Size (Scaled) = 101.41µm X 101.41µm
Bit Depth = 16 Bit
Image Center Position = X: 72.64mm, Y: 50.012mm
Stage Position = X: 72.64mm, Y: 50.012mm
ROI Center Offset = X: 0.00µm, Y: 0.00µm

488 -

Reflector = None
Contrast Method = Fluorescence
Pinhole = 1.00AU / 45µm
LMS MBS = MBS 405 + 488 + 561 + 640(T10/R90)
Laser Wavelength = 488nm @0.2%
Laser Blanking = Enabled
Scan Mode = Frame
Scan Zoom = 1.0
Rotation = 0°
Sampling = 1.00
Pixel Time = 0.65µs
Frame Time = 8.67s
LSM Scan Speed = 8
Scan Direction = Bidirectional
Line Step = 1
Averaging - 8
Excitation Wavelength = 493
Emission Wavelength = 517
Effective NA = 1.4
Detection Wavelength = 496 - 575
Imaging Device = Airyscan
Detector Type = GaAsp-PMT
Detector Gain = 750 V
Detector Offset = 0
Detector Digital Gain = 1.0

594 -

Reflector = None
Contrast Method = Fluorescence
Pinhole = 1.00AU / 54µm
LMS MBS = MBS 405 + 488 + 561 + 640(T10/R90)
Laser Wavelength = 561nm @0.2%
Laser Blanking = Enabled
Scan Mode = Frame
Scan Zoom = 1.0
Rotation = 0°
Sampling = 1.00
Pixel Time = 0.65µs
Frame Time = 8.67s
LSM Scan Speed = 8
Scan Direction = Bidirectional
Line Step = 1
Averaging - 8
Excitation Wavelength = 590
Emission Wavelength = 618
Effective NA = 1.4
Detection Wavelength = 590 - 700
Imaging Device = GaAsp-PMT 2
Detector Type = GaAsp-PMT
Detector Gain = 650 V
Detector Offset = 0
Detector Digital Gain = 1.0

Post Processing

Split Gallery, Frame 1 Identifyn® 594, Frame 2 Identifyn® 488, Frame 3 Merged.

Image Corrections

None

Image by J.K. Bennett

Copyright 2026 GMD12 LLC.

SOURCE PROVENANCE

Catalog	SA-000021
Stage	Confocal
Instrument	ZEISS LSM 900
Microscope configuration	Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:

SOURCE PROVENANCE

Structured source metadata values	39
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	1D668E4B7BC356A95517D4B418EBEBE8DBEC0B749B31A223CAF184F524D39ACB
Method PDF SHA-256	E5D86DEEC0E47666D500C45634F188B3CF1F25CAA15AF62576F71D361E5845F5

ORIGINAL IMAGE EVIDENCE / AIRYSCAN



EVIDENCE TIER	Airyscan
INSTRUMENT	ZEISS LSM 900
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488; 594
METADATA VALUES	46

STRUCTURED ACQUISITION METADATA / AIRYSCAN

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:
Objective	with a Plan-Apochromat 63X/1.40 oil DIC M27
Z-Stack	160 Slices (6.36µm)
Channels	2
Scaling (Per Pixel)	0.04252 µm X 0.04252 µm X 0.04000 µm
Image Size (Pixels)	1535 X 1348
Image Size (Scaled)	65.27µm X 57.32µm
Bit Depth	16 Bit
Image Center Position	X: 75.65mm, Y: 49.30mm
Stage Position	X: 75.65mm, Y: 49.30mm
ROI Center Offset	X: 0.00µm, Y: 0.00µm
Reflector	None
Contrast Method	Fluorescence
Pinhole	5.31AU / 236µm | 5.00AU / 272µm
LMS MBS	MBS 405 + 488 + 561 + 640(T10/R90)
Laser Wavelength	488nm @0.3% | 561nm @0.5%
Laser Blanking	Enabled
Scan Mode	Frame
Scan Zoom	1.5
Rotation	0°
Sampling	2.00
Pixel Time	0.66µs
Frame Time	3.94s
LSM Scan Speed	7
Scan Direction	Bidirectional
Line Step	1
Excitation Wavelength	493 | 590
Emission Wavelength	517 | 618
Effective NA	1.4
Detection Wavelength	495 - 569 | 598 - 700
Imaging Device	Airyscan
Detector Type	GaAsp-PMT
Detector Gain	850 V | 800 V

FIELD	SOURCE-DOCUMENTED VALUE
Detector Offset	0
Detector Digital Gain	1.0
AiryScan Mode	Super Resolution 9.1 (3D, Auto) | Super Resolution 8.3 (3D, Auto)
3D Volume Projection	Precise
Appearance	Threshold 6% (594), Threshold 6% (488), Ramp 94% (594), Ramp 94% (488), Maximum 100%
Background	Black
Light	Brightness100%, Azimuth 80°, Elongation -130°, Enabled Directional Lighting, Enabled Tone Mapping
Projection	View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

Identifyn® SRM Alexa Fluor™ 594 Affinity Purified Donkey Anti-Rabbit F(ab')□ (H + L)
Cat ID# - SA 000054

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher, Fish Serum Blocker cat#37527X3. The first primary antibody, Mitochondria Antibody (MA5-12017), was incubated overnight at 4°C at a dilution of 1:100. Cells were washed 3X in PBS. Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000.

The second primary antibody followed this, Actin Recombinant Rabbit Monoclonal Antibody (JF47-01), which was incubated for 1 hour at room temperature at a dilution of 1:250. Cells were washed 3X in PBS. Identifyn® 594 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:

Z Stack - (3D)

Z-Stack = 160 Slices (6.36µm)

Channels = 2
Scaling (Per Pixel) = 42.52nm X 42.52nmµm X 40.00nm
Image Size (Pixels) = 1535 X 1348
Image Size (Scaled) = 65.27µm X 57.32µm
Bit Depth = 16 Bit
Image Center Position = X: 75.65mm, Y: 49.30mm
Stage Position = X: 75.65mm, Y: 49.30mm
ROI Center Offset = X: 0.00µm, Y: 0.00µm

488 -

Reflector = None
Contrast Method = Fluorescence
Pinhole = 5.31AU / 236µm
LMS MBS = MBS 405 + 488 + 561 + 640(T10/R90)
Laser Wavelength = 488nm @0.3%
Laser Blanking = Enabled
Scan Mode = Frame
Scan Zoom = 1.5
Rotation = 0°
Sampling = 2.00
Pixel Time = 0.66µs
Frame Time = 3.94s
LSM Scan Speed = 7
Scan Direction = Bidirectional
Line Step = 1
Averaging - 2
Excitation Wavelength = 493
Emission Wavelength = 517
Effective NA = 1.4
Detection Wavelength = 495 - 569
Imaging Device = Airyscan
Detector Type = GaAsp-PMT
Detector Gain = 850 V
Detector Offset = 0
Detector Digital Gain = 1.0
AiryScan Mode = Super Resolution 9.1 (3D, Auto)

594 -

Reflector = None
Contrast Method = Fluorescence
Pinhole = 5.00AU / 272µm
LMS MBS = MBS 405 + 488 + 561 + 640(T10/R90)
Laser Wavelength = 561nm @0.5%
Laser Blanking = Enabled
Scan Mode = Frame
Scan Zoom = 1.5
Rotation = 0°
Sampling = 2.00
Pixel Time = 0.66µs
Frame Time = 3.94s
LSM Scan Speed = 7
Scan Direction = Bidirectional
Line Step = 1
Averaging - 2
Excitation Wavelength = 590
Emission Wavelength = 618
Effective NA = 1.4
Detection Wavelength = 598 - 700
Imaging Device = Airyscan
Detector Type = GaAsp-PMT
Detector Gain = 800 V
Detector Offset = 0
Detector Digital Gain = 1.0
AiryScan Mode = Super Resolution 8.3 (3D, Auto)

3D Image Processing

3D Volume Projection = Precise
Appearance = Threshold 6% (594), Threshold 6% (488), Ramp 94% (594), Ramp 94% (488), Maximum 100%
Background = Black
Light = Brightness100%, Azimuth 80°, Elongation -130°, Enabled Directional Lighting, Enabled Tone Mapping
Projection = View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

Image Corrections

None

Image by J.K. Bennett

SOURCE PROVENANCE

Catalog	SA-000021

SOURCE PROVENANCE

Stage	Airyscan
Instrument	ZEISS LSM 900
Microscope configuration	Zeiss LSM 900 Confocal, Axio Observer.Z1/7, using with a Plan-Apochromat 63X/1.40 oil DIC M27 was used with the following parameters for each dye:
Structured source metadata values	46
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	0104FFDB16D12BCE9471DABAA3C96326390D35BD7DB9DA9DAEFA37383E270E49
Method PDF SHA-256	664ACB37FD244D3155681F1ECEAC87F315B3E9F135EE140BE1ACA190E4FD5B40

ORIGINAL IMAGE EVIDENCE / SIM



EVIDENCE TIER	SIM
INSTRUMENT	ZEISS Elyra 7
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488; 594
METADATA VALUES	57

STRUCTURED ACQUISITION METADATA / SIM

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:
Objective	Plan-Apochromat 63X/1.46 oil Korr M27
Filters	-2147483648
Beam Splitter 488	Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Dual Camera Beam Splitter	SBS LP 560
Beam Splitter 594	Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Calibration Marker Positions	Marker 1: 0.00µm, 0.00µm, Marker 2: 0.00µm, 0.00µm, Marker 3 0.00µm, 0.00µm
Z-Stack	35Slices (2.04µm)
Channels	2
Scaling (Per Pixel)	0.03130 µm X 0.03130 µm X 0.06000 µm
Image Size (Pixels)	2560 X 2560
Image Size (Scaled)	80.14µm X 80.14µm
Bit Depth	16 Bit
Image Center Position	X: 3.07mm, Y: -768.70µm
Stage Position	X: 3.07mm, Y: -768.70µm
ROI Center Offset	X: 0.00µm, Y: 0.00µm
Contrast Method	Fluorescence
Laser Wavelength	488nm @3.50% | 561nm @4.0%
Scan Mode	Other
Scan Zoom	1.0
Rotation	0º
Frame Time	1.00s
Scan Direction	Unidirectional
Line Step	N/A
Averaging	8
Excitation Wavelength	488 | 561
Emission Wavelength	655 | 515
Effective NA	1.46
Detection Wavelength	655-1000 | N/A
Imaging Device	Edge 4.2 SIM
Exposure Time	65ms | 60ms
Depth of Focus	0.93µm | 0.73µm
Binning Mode	1x1

FIELD	SOURCE-DOCUMENTED VALUE
Detector Type	Camera
Depth Offset	0
Detector Digital Gain	0.00
Channel	488.000000 | 561.000000
Grating Period (µm)	27.500000
Processing	3D
Auto Sharpness	Yes
Sharpness	11.919940 | 10.621737
Detrend	No
Sectioning	100.000000
Baseline	Yes
Scale to Raw Image	No
Experimental PSF	No
3D Maximum Projection	Precise
Appearance	Threshold 6% (594), Threshold 8% (488), Ramp 0% (594), Ramp 0% (488), Maximum 100% both channels
Background	Black
Light	Brightness 100%, Enabled Tone Mapping
Projection	View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

Identifyn® SRM Alexa Fluor™ 594 Affinity Purified Donkey Anti-Rabbit F(ab')□ (H + L)
Cat ID# - SA 000054

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher, Fish Serum Blocker cat#37527X3. The first primary antibody, Mitochondria Antibody (MA5-12017), was incubated overnight at 4°C at a dilution of 1:100. Cells were washed 3X in PBS. Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000.

The second primary antibody followed this, Actin Recombinant Rabbit Monoclonal Antibody (JF47-01), which was incubated for 1 hour at room temperature at a dilution of 1:250. Cells were washed 3X in PBS. Identifyn® 594 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:

Filters = -2147483648

Beam Splitter 488 = Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Dual Camera Beam Splitter = SBS LP 560

Beam Splitter 594 = Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Dual Camera Beam Splitter = SBS LP 560

Calibration Marker Positions: Marker 1: 0.00µm, 0.00µm, Marker 2: 0.00µm, 0.00µm, Marker 3 0.00µm, 0.00µm

Z Stack - (3D)

Z-Stack = 35Slices (2.04µm)

Channels = 2
Scaling (Per Pixel) = 31.30nm X 31.30nm X 60.00nm
Image Size (Pixels) = 2560 X 2560
Image Size (Scaled) = 80.14µm X 80.14µm
Bit Depth = 16 Bit
Image Center Position = X: 3.07mm, Y: -768.70µm
Stage Position = X: 3.07mm, Y: -768.70µm
ROI Center Offset = X: 0.00µm, Y: 0.00µm

488 -

Contrast Method = Fluorescence
Laser Wavelength = 488nm @3.50%
Scan Mode = Other
Scan Zoom = 1.0
Rotation = 0º
Frame Time = 1.00s
Scan Direction = Unidirectional
Line Step = N/A
Averaging = 8
Excitation Wavelength = 488
Emission Wavelength = 655
Effective NA = 1.46
Detection Wavelength = 655-1000
Imaging Device = Edge 4.2 SIM
Exposure Time = 65ms
Depth of Focus = 0.93µm
Binning Mode = 1x1
Detector Type = Camera
Depth Offset = 0
Detector Digital Gain = 0.00

Post Processing - SIM Processing

Channel = 488.000000
Grating Period (µm): 27.500000

SIM

Processing: 3D
Auto Sharpness: Yes
Sharpness: 11.919940
Detrend: No
Sectioning: 100.000000
Baseline: Yes
Scale to Raw Image: No
Experimental PSF: No

594 -

Contrast Method = Fluorescence
Laser Wavelength = 561nm @4.0%
Scan Mode = Other
Scan Zoom = 1.0
Rotation = 0º
Frame Time = 1.00s
Scan Direction = Unidirectional
Line Step = N/A
Averaging = 8
Excitation Wavelength = 561
Emission Wavelength = 515
Detection Wavelength = N/A
Effective NA = 1.46
Imaging Device = Edge 4.2 SIM
Exposure Time = 60ms
Depth of Focus = 0.73µm
Binning Mode = 1x1
Detector Type = Camera
Depth Offset = 0
Detector Digital Gain = 0.00

Post Processing - SIM Processing

Channel = 561.000000
Grating Period (µm): 27.500000

SIM

Processing: 3D
Auto Sharpness: Yes
Sharpness: 10.621737
Detrend: No
Sectioning: 100.000000
Baseline: Yes
Scale to Raw Image: No
Experimental PSF: No

3D Image Processing

3D Maximum Projection = Precise
Appearance = Threshold 6% (594), Threshold 8% (488), Ramp 0% (594), Ramp 0% (488), Maximum 100% both channels
Background = Black
Light = Brightness 100%, Enabled Tone Mapping
Projection = View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

Image Corrections

None

Image by P. Raut

SOURCE PROVENANCE

Catalog	SA-000021
Stage	SIM
Instrument	ZEISS Elyra 7
Microscope configuration	Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:
Structured source metadata values	57
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	1AE6DFA6079D4D5326A6113C60ED6E22C3EEE0CD2D5B2D5791997E6B6C5C9B49
Method PDF SHA-256	74247839B24DF6A34FA245222DE2B00DB074F7ED0087602D55BC4BE369E4F85A

ORIGINAL IMAGE EVIDENCE / SIM²



EVIDENCE TIER	SIM²
INSTRUMENT	ZEISS Elyra 7
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488; 594
METADATA VALUES	62

STRUCTURED ACQUISITION METADATA / SIM²

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:
Objective	Plan-Apochromat 63X/1.46 oil Korr M27
Filters	-2147483648
Beam Splitter 594	Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Dual Camera Beam Splitter	SBS LP 560
Beam Splitter 488	Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Calibration Marker Positions	Marker 1: 0.00µm, 0.00µm, Marker 2: 0.00µm, 0.00µm, Marker 3 0.00µm, 0.00µm
Z-Stack	26Slices (1.5µm)
Channels	2
Scaling (Per Pixel)	0.01565 µm X 0.01565 µm X 0.06000 µm
Image Size (Pixels)	4734 X 2971 | 312 X 312
Image Size (Scaled)	74.09µm X 46.50µm | 4.88µm X 4.88µm
Bit Depth	16 Bit
Image Center Position	X: 2.63mm, Y: -1.50mm
Stage Position	X: 2.63mm, Y: -1.50mm
ROI Center Offset	X: 0.00µm, Y: 0.00µm
Contrast Method	Fluorescence
Laser Wavelength	488nm @3.50% | 561nm @4.0%
Scan Mode	Other
Scan Zoom	1.0
Rotation	0º
Frame Time	1.0s
Scan Direction	Unidirectional
Line Step	N/A
Averaging	8
Excitation Wavelength	488 | 561
Emission Wavelength	655 | 515
Effective NA	1.46
Imaging Device	Edge 4.2 SIM
Exposure Time	65ms | 60ms
Depth of Focus	0.93µm | 0.73µm
Binning Mode	1x1
Detector Type	Camera

FIELD	SOURCE-DOCUMENTED VALUE
Depth Offset	0
Detector Digital Gain	0.00
Channel	488.000000 | 561.000000
Grating Period (µm)	27.500000
Processing	3D
Input SNR	Medium
Iterations	16
Regularization Weight	0.065000
Processing Sampling	4
Output Sampling	4
Filter	Median
Detrend	No
Sectioning	100.000000
Baseline	Yes
Scale to Raw Image	No
Experimental PSF	No
3D Maximum Projection	Precise
Appearance	Threshold 4% (594), Threshold 2% (488) Ramp 0% (594), Ramp 0% (488), Maximum 100% both channels | Threshold 2% (594), Threshold 2% (488) Ramp 98% (594), Ramp 98% (488), Maximum 100% both channels
Background	Black
Light	Brightness 100%, Enabled Tone Mapping | Brightness 100%, Azimuth 80°, Elongation -130°, Enabled Directional Lighting, Enabled Tone Mapping
Projection	View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse F(ab')□ (H + L)
Cat ID# - SA 000021

Identifyn® SRM Alexa Fluor™ 594 Affinity Purified Donkey Anti-Rabbit F(ab')□ (H + L)
Cat ID# - SA 000054

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher, Fish Serum Blocker cat#37527X3. The first primary antibody, Mitochondria Antibody (MA5-12017), was incubated overnight at 4°C at a dilution of 1:100. Cells were washed 3X in PBS. Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000.

The second primary antibody followed this, Actin Recombinant Rabbit Monoclonal Antibody (JF47-01), which was incubated for 1 hour at room temperature at a dilution of 1:250. Cells were washed 3X in PBS. Identifyn® 594 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:

Filters = -2147483648

Beam Splitter 594 = Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Dual Camera Beam Splitter = SBS LP 560

Beam Splitter 488 = Main Beam Splitter Non-Descanned: LBF 405/488/561/642
Dual Camera Beam Splitter = SBS LP 560

Calibration Marker Positions: Marker 1: 0.00µm, 0.00µm, Marker 2: 0.00µm, 0.00µm, Marker 3 0.00µm, 0.00µm

Z Stack - (3D) - Main Image

Z-Stack = 26Slices (1.5µm)

Channels = 2
Scaling (Per Pixel) = 15.65nm X 15.65nm X 60.00nm
Image Size (Pixels) = 4734 X 2971
Image Size (Scaled) = 74.09µm X 46.50µm
Bit Depth = 16 Bit
Image Center Position = X: 2.63mm, Y: -1.50mm
Stage Position = X: 2.63mm, Y: -1.50mm
ROI Center Offset = X: 0.00µm, Y: 0.00µm

Z Stack - (3D) - Zoomed/Inset

Z-Stack = 26Slices (1.5µm)

Channels = 2
Scaling (Per Pixel) = 15.65nm X 15.65nm X 60.00nm
Image Size (Pixels) = 312 X 312
Image Size (Scaled) = 4.88µm X 4.88µm
Bit Depth = 16 Bit
Image Center Position = X: 2.63mm, Y: -1.50mm
Stage Position = X: 2.63mm, Y: -1.50mm
ROI Center Offset = X: 0.00µm, Y: 0.00µm

488 -

Contrast Method = Fluorescence
Laser Wavelength = 488nm @3.50%
Scan Mode = Other
Scan Zoom = 1.0
Rotation = 0º
Frame Time = 1.0s
Scan Direction = Unidirectional
Line Step = N/A
Averaging = 8
Excitation Wavelength = 488
Emission Wavelength = 655
Effective NA = 1.46
Imaging Device = Edge 4.2 SIM
Exposure Time = 65ms
Depth of Focus = 0.93µm
Binning Mode = 1x1
Detector Type = Camera
Depth Offset = 0
Detector Digital Gain = 0.00

Post Processing - SIM2 Processing

Channel = 488.000000
Grating Period (µm): 27.500000

SIM2
Processing: 3D
Input SNR: Medium
Iterations: 16
Regularization Weight: 0.065000
Processing Sampling: 4
Output Sampling: 4
Filter: Median
Detrend: No
Sectioning: 100.000000
Baseline: Yes
Scale to Raw Image: No
Experimental PSF: No

594 -

Contrast Method = Fluorescence
Laser Wavelength = 561nm @4.0%
Scan Mode = Other
Scan Zoom = 1.0
Rotation = 0º
Frame Time = 1.0s
Scan Direction = Unidirectional
Line Step = N/A
Averaging = 8
Excitation Wavelength = 561
Emission Wavelength = 515
Effective NA = 1.46
Imaging Device = Edge 4.2 SIM
Exposure Time = 60ms
Depth of Focus = 0.73µm
Binning Mode = 1x1
Detector Type = Camera
Depth Offset = 0
Detector Digital Gain = 0.00

Post Processing - SIM2 Processing

Channel = 561.000000
Grating Period (µm): 27.500000

SIM2
Processing: 3D
Input SNR: Medium
Iterations: 16
Regularization Weight: 0.065000
Processing Sampling: 4
Output Sampling: 4
Filter: Median
Detrend: No
Sectioning: 100.000000
Baseline: Yes
Scale to Raw Image: No
Experimental PSF: No

3D Image Processing - Main Image

3D Maximum Projection = Precise
Appearance = Threshold 4% (594), Threshold 2% (488) Ramp 0% (594), Ramp 0% (488), Maximum 100% both channels
Background = Black
Light = Brightness 100%, Enabled Tone Mapping
Projection = View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

3D Image Processing - Zoomed/Inset

3D Maximum Projection = Precise
Appearance = Threshold 2% (594), Threshold 2% (488) Ramp 98% (594), Ramp 98% (488), Maximum 100% both channels
Background = Black
Light = Brightness 100%, Azimuth 80°, Elongation -130°, Enabled Directional Lighting, Enabled Tone Mapping
Projection = View Angle 35°, Scale Z 100%, Reverse Z-Axis NOT Selected, Stereo Anaglyph NOT Selected, Camera Separation 5% (deselected), Parallax Shift 0% (deselected)

Image Corrections

None

Image by P. Raut

SOURCE PROVENANCE

Catalog	SA-000021

SOURCE PROVENANCE

Stage	SIM²
Instrument	ZEISS Elyra 7
Microscope configuration	Zeiss Elyra 7, Andor1, AxioObserver, using Plan-Apochromat 63X/1.46 oil Korr M27 was used with the following parameters for each dye:
Structured source metadata values	62
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	DD6859DA65D27704C2E607BD59229C0599481AC6489C84A01D0D2D8C2D203BE0
Method PDF SHA-256	8BB1A051AC4F6EE133870BB9C7E19FFD35144E211F5786DFF4C94DC2587AA107

ORIGINAL IMAGE EVIDENCE / SINGLE-MOLECULE STORM



EVIDENCE TIER	Single-molecule STORM
INSTRUMENT	Bruker Vutara VXL
CELL MODEL	HeLa
DETECTED DYES	488
METADATA VALUES	71

STRUCTURED ACQUISITION METADATA / SINGLE-MOLECULE STORM

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Bruker Vutara VXL Biplane Single Molecule Microscope, using an Olympus 60X/1.30 silicon oil objective, was used with the following parameters:
Objective	Olympus 60X/1.30 silicon oil objective
Number of Probes	1
Number of Simultaneous Probes	1
Number of Reference Probes	1
Camera	Both Cameras
Color Channel	Both Channels
Probe Capture Mode	Interleaved
Z -Stack Capture Mode	Interleaved
Multiple Location Capture	Not selected
Post Record Reference	Not Selected
Fluidics	Not Selected
Autofocus Drift Correction	On
Biplane Module	635nm Dichroic
Objective	Silicon Selected
Stage Insert	Rectangular Selected
Course Focus Position	8-Well Chamber Selected
Dichroic	SuperRes
PSF	(Red, Orange, Green)
Heater	Current: 26
Recording Vertical Field of View	100%
Widefield Camera Exposure Time	100ms | 20ms
Super Resolution Camera Exposure Time	20ms
Widefield Camera Binning	1
Per-Probe Exposure Time	Selected
Predefined Probe	First probe: 488 nm (635 nm Dichroic); First reference probe: 488 nm (635 nm Dichroic)
Photoactivation/Imaging Laser	0% Selected
Exposure Time	20ms Selected
Record Workflow	Selected Options
View Mode	Widefield 200µm
Capture Mode	Live
Piezo Current	179 µm

FIELD	SOURCE-DOCUMENTED VALUE
Widefield 488nm	0.1X Neutral Density Filter: Selected, @0.02% Laser Input
Reference Probe 1 Laser control (488 nm (635 nm Dichroic)	Not Selected
Widefield camera Exposure Time	100ms
Widefield Laser	"Widefield 488nm" was deselected, and "calibrate" under Autofocus was selected.
Current Intensity	4.5
Calibration Value	-5084 nm/unit
Autofocus Mode	Pre-Capture
SuperRes	50µm is Selected
Piezo Record	Begin: 178; End: 178
SuperRes 488nm	0.1X Neutral Density Filter: Selected, @0.01% Laser Input
Record Workflow - Record Tab	Capture Parameters
Frames	Probe 1: 20000
Z Positions	1 using Steps: 0.2µm (200nm)
Cycles	1
Time Points	1
Pause Between Cycles	Not Selected
Automatic Photoactivation Laser Control	Not Selected
Laser Power	9%
Cutout Width	16px
Max Frame Accumulation	1 frame
Max Accumulation Offset	1px
Fiducial at Max Accumulation	Not selected
Max Particles Per Frame	No Max
Background SLPE Removal	Not Selected
Cutout/Background Removal	Not Selected
Preview Cutout Positions	Selected
Probe1	BG threshold: 7
Sizing Mode	Constant
Particle size	25nm
Color by	Constant
Color Table	Single
Opacity Mode	Constant; Opacity 1: 7
Front-to-back Attenuation	Not Selected
Method	Particle (direct)
Time Intervals	5 Intervals

FIELD	SOURCE-DOCUMENTED VALUE
Pixel Size	20nm
Smoothing	8.00
PSF Z offset	-450 to +450
Radial precision	35
Axial precision	70

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse, F(ab')□ (H + L)
Cat ID# - SA 000021

Hela cells were grown on µ-Slide 8 Well high Glass Bottom Ibidi chamber, followed by -20°C methanol fixation and blocked with Thermo Fisher Fish Serum Blocker cat#37527X3. The primary antibody, Alpha Tubulin Monoclonal Antibody (DM1A), was incubated for 1 hour at room temperature at a dilution of 1:100. Cells were washed 3X in Identifyn's™ proprietary Wash Buffer, and Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:100. Cells were washed 5X in Identifyn's™ proprietary Wash Buffer followed by the addition of Identifyn's™ proprietary STORM Buffer to the sample chamber enabling image acquisition.

Microscope

Bruker Vutara VXL Biplane Single Molecule Microscope, using an Olympus 60X/1.30 silicon oil objective, was used with the following parameters:

Calibration -

Calibration follows Identifyn's Calibration Standard Operating Procedure, which can be found here.

https://identifyn.com/storm-calibration

Image Acquisition -

SRX Software - Under the 'Single Molecule Localization' tab, 'Record' is selected and contains the following workflow tabs and parameters.

Configuration Tab - Capture Configuration

Number of Probes: 1
Number of Simultaneous Probes: 1
Number of Reference Probes: 1
Camera: Both Cameras
Color Channel: Both Channels
Probe Capture Mode: Interleaved
Z -Stack Capture Mode: Interleaved
Multiple Location Capture: Not selected
Post Record Reference: Not Selected
Fluidics: Not Selected
Autofocus Drift Correction: On

Microscope configuration

Biplane Module: 635nm Dichroic
Objective: Silicon Selected
Stage Insert: Rectangular Selected
Course Focus Position: 8-Well Chamber Selected
Dichroic: SuperRes
PSF: (Red, Orange, Green)
Heater: Current: 26

Camera Configuration

Recording Vertical Field of View: 100%
Widefield Camera Exposure Time: 100ms
Super Resolution Camera Exposure Time: 20ms
Widefield Camera Binning: 1
Per-Probe Exposure Time: Selected

Probe Configuration

Predefined Probe: First probe: 488 nm (635 nm Dichroic); First reference probe: 488 nm (635 nm Dichroic)
Photoactivation/Imaging Laser: 0% Selected
Exposure Time: 20ms Selected

Experiment Configuration

Record Workflow: Selected Options
View Mode: Widefield 200µm
Capture Mode: Live

Stage and Piezo Selection

Examination of Sample

Cell(s) or Cellular Structure Localization - The red-bordered box identifies the area of the SuperRes view and defines a suitable cell(s) or cellular structure. The cell(s) or structures are optimally focused.

Piezo Current: 179 µm

Probe 1 Laser control (488 nm (635 nm Dichroic)

Widefield 488nm: 0.1X Neutral Density Filter: Selected, @0.02% Laser Input
Reference Probe 1 Laser control (488 nm (635 nm Dichroic): Not Selected

Advanced Tools Option

Widefield camera Exposure Time: 100ms
Default values are maintained unless otherwise noted.

Widefield Laser: "Widefield 488nm" was deselected, and "calibrate" under Autofocus was selected.

Autofocus - Calibration Values

Current Position" -0.979
Current Intensity: 4.5
Calibration Value: -5084 nm/unit
Autofocus Mode: Pre-Capture

View Mode

SuperRes: 50µm is Selected

Capture Mode

Live

Stage and Piezo Selection

Cell(s) or Cellular Structure Localization - Positioning of the target Cell or Subsection of the Target Cell is optimized and optimally focused.

Piezo Record: Begin: 178; End: 178

Probe 1 Laser control (488 nm (635 nm Dichroic)

SuperRes 488nm: 0.1X Neutral Density Filter: Selected, @0.01% Laser Input

Reference Probe 1 Laser control (488 nm (635 nm Dichroic)

Not Selected

Advanced Tools

Widefield Camera Exposure Time: 20ms
Default values are maintained unless otherwise noted.

Record Workflow - Record Tab: Capture Parameters

Frames: Probe 1: 20000
Z Positions: 1 using Steps: 0.2µm (200nm)
Cycles: 1
Time Points: 1
Pause Between Cycles: Not Selected
Automatic Photoactivation Laser Control: Not Selected

Probe 1 Laser control (488 nm (635 nm Dichroic)

Laser Power: 9%

Expert Configuration Options - The Localization Tab

Beyond the values shown, all other parameters were default values.

Localization

Cutout Width: 16px
Max Frame Accumulation: 1 frame
Max Accumulation Offset: 1px
Fiducial at Max Accumulation: Not selected
Max Particles Per Frame: No Max
Background SLPE Removal: Not Selected
Cutout/Background Removal: Not Selected
Preview Cutout Positions: Selected

Record Workflow - Localization Tab

Data Collection

The folder is created automatically with the localization file for this image acquisition.

Recording Summary

All parameters were carried over from previous tabs and are kept except:

Probe1: BG threshold: 7

Region of Interest -

Not selected

Display

Was deployed to inspect if the BG threshold applied chooses localizations in an expected manner

Record Workflow - View Tab

Region of Interest

Default values

Visualization Parameters

Point Splatting
Sizing Mode: Constant
Particle size: 25nm
Color by: Constant
Color Table: Single
Opacity Mode: Constant; Opacity 1: 7
Front-to-back Attenuation: Not Selected

Post Acquisition analysis - View Tab

Drift Correction - Computed with the following parameters:

Method: Particle (direct)
Time Intervals: 5 Intervals
Pixel Size: 20nm
Smoothing: 8.00

Advanced Particle Filters - Applied to the drift-corrected data with the following parameters

PSF Z offset: -450 to +450
Radial precision: 35
Axial precision: 70

The other parameters were not applied.

Statistical analysis - N/A

Image by S. Thakur

SOURCE PROVENANCE

Catalog	SA-000021
Stage	Single-molecule STORM
Instrument	Bruker Vutara VXL
Microscope configuration	Bruker Vutara VXL Biplane Single Molecule Microscope, using an Olympus 60X/1.30 silicon oil objective, was used with the following parameters:
Structured source metadata values	71
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa
Image SHA-256	DE97F675ED3E53204FD5777344B129C0C36130DF070A765C48744EC0B9578F12
Method PDF SHA-256	8D688A16C7D35185620744158FC618A0AD774D19FD554848436EFD099A790228

ORIGINAL IMAGE EVIDENCE / CONTROL



EVIDENCE TIER	Control
INSTRUMENT	ZEISS Axio Imager.M1; ZEISS Axio Observer.Z1/7 Apotome
CELL MODEL	HeLa
DETECTED DYES	405/DAPI; 488
METADATA VALUES	30

STRUCTURED ACQUISITION METADATA / CONTROL

The values below were deterministically extracted from the preserved Identifyn method PDF text. They remain source-documented metadata, not new BENNETT measurements.

FIELD	SOURCE-DOCUMENTED VALUE
Platform / configuration	Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:
Objective	EC Plan-Neofluar 63X/1.25 oil M27
Channels	2
Scaling (Per Pixel)	0.143µm X 0.143µm
Image Size (Pixels)	1600 X 1104
Image Size (Scaled)	228.57µm X 157.71µm
Bit Depth	14 Bit
Image Center Position	X: 46.29mm, Y: 52.38mm
ROI Center Offset	X: 46.29µm, Y: 52.38µm
Reflector	38 HE eGFP | 49 DAPI
Beam Splitter	495 | 395
Filter Excitation Wavelength	450 - 490 | 335-383
Filter Emission Wavelength	500 -550 | 420-470
Contrast Method	Fluorescence
Light source	X-Cite Xylis Lamp Intensity @25% | X-Cite Xylis Lamp Intensity @10%
Exposure Time	200ms | 30ms
Excitation Wavelength	493 | 353
Emission Wavelength	517 | 465
Effective NA	1.25
Imaging Device	Axiocam 807 mono
Camera Adapter	1X
Depth of Focus	1.0µm | 0.90µm
Binning Mode	2,2

SOURCE METHOD

The text below is carried from the preserved Identifyn source record. Typography and pagination have been normalized for this preservation dossier; scientific wording has not been editorially rewritten.

Identifyn Standards for Optical Microscopy Methods™

Identifyn® SRM Alexa Fluor™ 488 Affinity Purified Rabbit Anti-Mouse, F(ab')□ (H + L)
Cat ID# - SA 000021

Hela Cells were grown on acid-etched #1.5H cover glass to ~50% confluency, followed by -20°C methanol fixation and blocked with Thermo Fisher Fish Serum Blocker cat#37527X3. In the absence of a primary antibody, the Identifyn® 488 secondary antibody was incubated at room temperature for 1 hour at a dilution of 1:1000. Cells were washed 5X in PBS and mounted using ProLong™ Diamond Antifade Mountant with DAPI cat#P36962.

Microscope

Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:

Single Plane (2D)

Channels = 2
Scaling (Per Pixel) = 0.143µm X 0.143µm
Image Size (Pixels) = 1600 X 1104
Image Size (Scaled) = 228.57µm X 157.71µm
Bit Depth = 14 Bit
Image Center Position = X: 46.29mm, Y: 52.38mm
ROI Center Offset = X: 46.29µm, Y: 52.38µm

488 -

Reflector = 38 HE eGFP
Beam Splitter = 495
Filter Excitation Wavelength = 450 - 490
Filter Emission Wavelength = 500 -550
Contrast Method = Fluorescence
Light source = X-Cite Xylis Lamp Intensity @25%
Exposure Time = 200ms
Excitation Wavelength = 493
Emission Wavelength = 517
Effective NA = 1.25
Imaging Device = Axiocam 807 mono
Camera Adapter = 1X
Depth of Focus = 1.0µm
Binning Mode = 2,2

DAPI -

Reflector = 49 DAPI
Beam Splitter = 395
Filter Excitation Wavelength = 335-383
Filter Emission Wavelength = 420-470
Contrast Method = Fluorescence
Light source = X-Cite Xylis Lamp Intensity @10%
Exposure Time = 30ms
Excitation Wavelength = 353
Emission Wavelength = 465
Effective NA = 1.25
Imaging Device = Axiocam 807 mono
Camera Adapter = 1X
Depth of Focus = 0.90µm
Binning Mode = 2,2

Post Processing -

None

Image Corrections

None

Control Summary -

In the absence of a primary antibody, Identifyn Standards for Optical Microscopy Methods™ were applied using standard staining methodology and image acquisition settings using the Zeiss Axio Imager.M1 Widefield. With no image correction, the image demonstrates that the secondary antibody, minus a primary target, had no off-target binding and no significant presence of free dye within the sample.

The data suggests the secondary antibody is highly specific and free of unconjugated (free) dye.

Image by J.K. Bennett

SOURCE PROVENANCE

Catalog	SA-000021
Stage	Control
Instrument	ZEISS Axio Imager.M1; ZEISS Axio Observer.Z1/7 Apotome
Microscope configuration	Zeiss Axio Observer.Z1/7 Apotome SIM, using an EC Plan-Neofluar 63X/1.25 oil M27, Axiocam 807, and X-Cite Xylis Lamp was used with the following parameters for each dye:
Structured source metadata values	30
Metadata source	PRESERVED_SOURCE_METHOD_PDF_TEXT
Cell model	HeLa

SOURCE PROVENANCE

Image SHA-256	708C6FF758CB0D595FBDF417E5B1AEC74AFD1ADBC27A80D499D31BEB65D321AE
Method PDF SHA-256	D0A53C497EAA6D6D9ED19E46F8D62D14284E3F3B6E3750BC2B8BD79F6A2356A8